

*Therapeutic drugs of pet animals*

## **\*Antibiotics\***

**oxytetracycline    1 cm /10 kg**

**sulfa dimidine 33.3%    0.2 gm / kg**

**1-Ampicillin vial (500 gm)**

(Epicocillin  $\frac{1}{2}$  gm)(Emox  $\frac{1}{2}$  gm)

1vial / 12 hr / I.M    3-5 days

**2-Ampicillin susp.:**

// 250    5 ml / 8 hr for 3-5 days

// 125    5 ml / 6 hr for 3-5 days

**3-E.mox susp. (amoxicillin)**

250    5 ml / 8 hr for 3-5 days

125    5 ml / 6 hr for 3-5 days

4-Erythromycin 200

(resp. Tract infection)

5 ml / 6 hr for 3-5 days

5-Velosef (½ gm), (Cefotax ½ gm)

1 vial / twice daily / I.M for 3-5 days

6-Flumox (½ gm)

1 vial / twice daily / I.M for 3-5 days

## **Antidiarrhoea**

**1-cidostin (susp) chlorumphenicol**

-enteritis , collibacilosis , salmonellosis

5 ml / 6 hr

**2-Kapect comp. (kaolin, sulfa , trimethopprime)**

20 ml / 12 hr

**3-Enterokin**

20 ml / 12 hr

**4-Flagyl (metronidazol) or (antinal) or (furazol) or  
(amrimzol)**

5 ml / 8 hr

**5-Rehydran N.pack ( rehydrozinc pack)**

one pack / 12 hr

## **\*digestant drugs \***

**\*Digestin syr    10 cm / 3 times / before meal ¼ hr**

**\*Neo-digestin syr    the same**

## **\*apptizer drug\***

**Vitaphos syr    5 cm / 3 times / before meal**

**Mosegor syr    the same**

## **\*antiemetic\***

**Primpran    amp**

**Plasil    amp                  1 amp / day / I.M**

**cortigen B6 amp**

## **\*laxative\***

**Sedalac (lactulose) syr**

**Lactulose (lactulose) syr    20 cm / 12 hr**

## \*antiflatulance\*

**Simethicone** syr 5 cm / 3 times / daily

**Flatidyl tab (disflatyl tab )** 5 tablet / orally / 3 times daily

## \*anthelmintics\*

Albendazol susp 5% 0.1 ml / kg

Ucimisole (levamisole) injection 10%

0.5 ml / 10 kg / S.C once

**\*Antiver susp**

**\*Fluvermol susp** 5 cm / 12 hr / for 3 days

**\*Fluver**

## \*Drugs for influenza & common cold\*

**Flurest syr**

**Noflu syr**

**Rhinomol syr**      **5 cm / 12 hr / 3-5 days**

**Rhinopront syr**

### \*antipyretic\*

\*cold application firstly then:

**Analgin**    **1-5 cm I/M**

**Novacid**    **1-5 cm I/M**

**\*paramol syr**

**\*paracetamol syr**      **5 cm / 8 hr**

**\*Cetal syr**

## **\*Expectorant & cough sedative\***

**Bronchophane syr**

**Expectyl syr**

**Toplexil syr**

**5 cm / 12 hr / 3-5 days**

**Bronchistal syr**

## **Mucolytics**

- **Bisolvon syr**
- **Mucosphylline**
- **Mucopront**
- **Bronchopront**

**5 cm/ 12 hr / 3-5 days**

## **Bronchodilator:**

- **Minophyline syr**
- **Salbovent syr**
- **Epicophyline syr**

**5 cm / 12 hr / 3-5 days**

## Antitussive for dry cough

- Tussilar syr
- Codilar syr
- Codipront syr                    5 cm / 12 hr / 3-5 days
- Coflin syr
- Codiphane syr

## Vitamines & general tonic

- Medivit syr
- Vitacal syr                    5 cm / 12 hr
- Multi sanostol syr
- Catosol                        0.5-5 cm (dog), 0.5-2.5 cm (cat)

## Anti inflammatory

- Dexamethosone injection  
0.1 – 1ml / daily

## Antitoxicisty

Atropin sulfste 1 %

## Topical antirhumatic & antiinflammatory

- Voltarin gel
- Olfen gel
- Profenid gel                          TWICE / DAILY
- Fostum gel

## Urinary antiseptic

- Coli urinal                          10 gm / 200 cm / twice daily
- Uricol

## Antihistaminic & antiallergic

- Allergyl syr
- Histalin syr                          5 cm / once / at night

## Topical drug for Scabies

- Benzonil emluge
- Benzail benzoate lotion twice/ daily
- Lice spray & lotion
- Eurax lotion

## Topical antifungal

- Canestan cream
- Mycostatin cream twice/ daily
- Daktacort cream

## Topical antihistaminic & antiallergic

- Betaderm cream
- Allergyl cream twice / daily
- Dexapanthen cream
- Diprosalic (betamethasone + salcylic acid )

## Topical antibiotic

- Terramycin oint. Twice / daily
- Garamycin oint.

## Topical antiseptic

- Betadine
- Savlon

## Eye infection

- Visin Ac eye drops 3 times / daily
- Prizolin eye drops  
  
(for relieve of redding)

### \*antibiotics:

- Isoptophenicol e.drop
- Isomephenicol e.drop 3 times / daily
- Isoptomaxitrol ( neomycin , dexamethasone)

\*\*Terramycin eye oint. At night

# Vaccination

## 1- Tetra valent vaccine (modified live)

- Canine distemper
- Adeno virus
- Para influenza
- Parvo virus
- N.B: (penta vaccine as previous + leptospira)

**At 1.5 – 2 month of age, I/M**

## 2- Rabies vaccine (inactivated vaccine)

at 3 month of age



6 month of age



anually